



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/715, 19/00	A1	(11) International Publication Number: WO 97/33913 (43) International Publication Date: 18 September 1997 (18.09.97)
<p>(21) International Application Number: PCT/US97/04043</p> <p>(22) International Filing Date: 12 March 1997 (12.03.97)</p> <p>(30) Priority Data: 60/013,345 13 March 1996 (13.03.96) US</p> <p>(71) Applicant: ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</p> <p>(72) Inventors: BAUMGARTNER, James, W.; 3504 Radcliffe Road, Thousand Oaks, CA 91360 (US). FARRAH, Theresa, M.; 718 16th Avenue, Seattle, WA 98127 (US). FOSTER, Donald, C.; 3002 N.E. 181st Street, Seattle, WA 98155 (US). GRANT, Francis, J.; 7714 37th Avenue N.E., Seattle, WA 98115 (US). O'HARA, Patrick, J.; 515 North 64th, Seattle, WA 98103 (US).</p> <p>(74) Agent: PARKER, Gary, E.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: CYTOKINE-RECEPTOR EXPRESSED IN TESTIS CELLS

(57) Abstract

Novel receptor polypeptides, polynucleotides encoding the polypeptides, and related compositions and methods are disclosed. The polypeptides comprise an extracellular domain of a cell-surface receptor that is expressed in testis cells. The polypeptides may be used within methods for detecting ligands that promote the proliferation and/or differentiation of testis cells, and may also be used in the development of male-specific contraceptives and infertility treatments.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Description

CYTOKINE-RECEPTOR EXPRESSED IN TESTIS CELLS

5

Background of the Invention

Proliferation and differentiation of cells of multicellular organisms are controlled by hormones and polypeptide growth factors. These diffusable molecules
10 allow cells to communicate with each other and act in concert to form cells and organs, and to repair and regenerate damaged tissue. Examples of hormones and growth factors include the steroid hormones (e.g. estrogen, testosterone), parathyroid hormone, follicle
15 stimulating hormone, the interleukins, platelet derived growth factor (PDGF), epidermal growth factor (EGF), granulocyte-macrophage colony stimulating factor (GM-CSF), erythropoietin (EPO) and calcitonin.

Hormones and growth factors influence cellular
20 metabolism by binding to receptors. Receptors may be integral membrane proteins that are linked to signalling pathways within the cell, such as second messenger systems. Other classes of receptors are soluble molecules, such as the transcription factors.

25 Of particular interest are receptors for cytokines, molecules that promote the proliferation and/or differentiation of cells. Examples of cytokines include erythropoietin (EPO), which stimulates the development of red blood cells; thrombopoietin (TPO), which stimulates
30 development of cells of the megakaryocyte lineage; and granulocyte-colony stimulating factor (G-CSF), which stimulates development of neutrophils. These cytokines are useful in restoring normal blood cell levels in patients suffering from anemia or receiving chemotherapy
35 for cancer. The demonstrated *in vivo* activities of these cytokines illustrates the enormous clinical potential of, and need for, other cytokines, cytokine agonists, and

cytokine antagonists. The present invention addresses this need by providing novel cytokine receptors and related compositions and methods.

5 Summary of the Invention

Within one aspect, the present invention provides an isolated polynucleotide encoding a ligand-binding receptor polypeptide. The polypeptide comprises a sequence of amino acids selected from the group consisting
10 of (a) residues 141 to 337 of SEQ ID NO:2; (b) allelic variants of (a); and (c) sequences that are at least 80% identical to (a) or (b). Within one embodiment, the polypeptide comprises residues 141 to 337 of SEQ ID NO:2 or SEQ ID NO:4. Within another embodiment, the
15 polypeptide encoded by the isolated polynucleotide further comprises a transmembrane domain. The transmembrane domain may comprise residues 340 to 363 of SEQ ID NO:2, or an allelic variant thereof. Within another embodiment, the polypeptide encoded by the isolated polynucleotide
20 further comprises an intracellular domain, such as an intracellular domain comprising residues 364 to 380 of SEQ ID NO:2, or an allelic variant thereof. Within further embodiments, the polynucleotide encodes a polypeptide that comprises residues 25 to 337, 1 to 337, or 1 to 380 of SEQ
25 ID NO:2 or SEQ ID NO:4. Within an additional embodiment, the polypeptide further comprises an affinity tag. Within a further embodiment, the polynucleotide is DNA.

Within a second aspect of the invention there is provided an expression vector comprising (a) a
30 transcription promoter; (b) a DNA segment encoding a secretory peptide and a ligand-binding receptor polypeptide, wherein the polypeptide comprises a sequence of amino acids selected from the group consisting of: (i) residues 141 to 337 of SEQ ID NO:2; (ii) allelic variants
35 of (i); and (iii) sequences that are at least 80% identical to (i) or (ii); and (c) a transcription terminator, wherein the promoter, DNA segment, and

terminator are operably linked. The ligand-binding receptor polypeptide may further comprise a transmembrane domain, or a transmembrane domain and an intracellular domain.

5 Within a third aspect of the invention there is provided a cultured eukaryotic cell into which has been introduced an expression vector as disclosed above, wherein said cell expresses a receptor polypeptide encoded by the DNA segment. Within one embodiment, the cell
10 further expresses a signalling subunit, such as a hematopoietic receptor β_c subunit. Within another embodiment, the cell is dependent upon an exogenously supplied hematopoietic growth factor for proliferation.

 Within a fourth aspect of the invention there is
15 provided an isolated polypeptide comprising a segment selected from the group consisting of (a) residues 141 to 337 of SEQ ID NO:2; (b) allelic variants of (a); and (c) sequences that are at least 80% identical to (a) or (b), wherein said polypeptide is substantially free of
20 transmembrane and intracellular domains ordinarily associated with hematopoietic receptors. Within one embodiment, the polypeptide further comprises an immunoglobulin F_c polypeptide. Within a another embodiment, the polypeptide further comprises an affinity
25 tag, such as polyhistidine, protein A, glutathione S transferase, or an immunoglobulin heavy chain constant region. Within a further embodiment, the polypeptide comprises residues 25-337 of SEQ ID NO:2, an allelic variant of SEQ ID NO:2, or a sequence that is at least 80%
30 identical to residues 25-337 of SEQ ID NO:2 or an allelic variant of SEQ ID NO:2.

 Within a further aspect of the invention there is provided a chimeric polypeptide consisting essentially of a first portion and a second portion joined by a
35 peptide bond. The first portion of the chimeric polypeptide consists essentially of a ligand binding domain of a receptor polypeptide selected from the group

consisting of (a) a receptor polypeptide as shown in SEQ ID NO:2; (b) allelic variants of SEQ ID NO:2; and (c) receptor polypeptides that are at least 80% identical to (a) or (b). The second portion of the chimeric polypeptide consists essentially of an affinity tag. Within one embodiment the affinity tag is an immunoglobulin F_C polypeptide. The invention also provides expression vectors encoding the chimeric polypeptides and host cells transfected to produce the chimeric polypeptides.

The invention also provides a method for detecting a ligand within a test sample, comprising contacting a test sample with a polypeptide as disclosed above, and detecting binding of the polypeptide to ligand in the sample. Within one embodiment the polypeptide further comprises transmembrane and intracellular domains. The polypeptide can be membrane bound within a cultured cell, wherein the detecting step comprises measuring a biological response in the cultured cell. Within another embodiment, the polypeptide is immobilized on a solid support.

Within an additional aspect of the invention there is provided an antibody that specifically binds to a polypeptide as disclosed above.

These and other aspects of the invention will become evident upon reference to the following detailed description and the attached drawing.

Brief Description of the Drawing

The Figure illustrates conserved structural features in cytokine receptors.

Detailed Description of the Invention

The term "allelic variant" is used herein to denote any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in

phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequence. The term allelic variant is also used herein to denote a protein encoded by an allelic variant of a gene.

The term "expression vector" is used to denote a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems.

"Operably linked", when referring to DNA segments, indicates that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in the promoter and proceeds through the coding segment to the terminator.

A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules.

The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter

sequences are commonly, but not always, found in the 5' non-coding regions of genes.

The term "receptor" is used herein to denote a cell-associated protein, or a polypeptide subunit of such a protein, that binds to a bioactive molecule (the "ligand") and mediates the effect of the ligand on the cell. Binding of ligand to receptor results in a conformational change in the receptor (and, in some cases, receptor multimerization, i.e., association of identical or different receptor subunits) that causes interactions between the effector domain(s) and other molecule(s) in the cell. These interactions in turn lead to alterations in the metabolism of the cell. Metabolic events that are linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, cell proliferation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids. The term "receptor polypeptide" is used to denote complete receptor polypeptide chains and portions thereof, including isolated functional domains (e.g., ligand-binding domains).

A "secretory signal sequence" is a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

A "soluble receptor" is a receptor polypeptide that is not bound to a cell membrane. Soluble receptors are most commonly ligand-binding receptor polypeptides that lack transmembrane and cytoplasmic domains. Soluble receptors can comprise additional amino acid residues, such as affinity tags that provide for purification of the polypeptide or provide sites for attachment of the

polypeptide to a substrate, or immunoglobulin constant region sequences. Many cell-surface receptors have naturally occurring, soluble counterparts that are produced by proteolysis or translated from alternatively spliced mRNAs. Receptor polypeptides are said to be substantially free of transmembrane and intracellular polypeptide segments when they lack sufficient portions of these segments to provide membrane anchoring or signal transduction, respectively.

The present invention is based in part upon the discovery of a novel DNA sequence that encodes a protein having the structure of a cytokine receptor, including the conserved WSXWS motif (SEQ ID NO:5). Analysis of the tissue distribution of the mRNA corresponding to this novel DNA showed that it was highly expressed in the testes, suggesting that the receptor mediates processes of progenitor cell growth and development, such as spermatogenesis. The receptor is also expressed at lower levels in pituitary. Subsequently, the receptor was shown to bind interleukin 13 (IL-13). The human cDNA was subsequently used to clone the orthologous receptor from Celebus macaque. The receptor has been designated "ZCyt2".

Cytokine receptors subunits are characterized by a multi-domain structure comprising a ligand-binding domain and an effector domain that is typically involved in signal transduction. Multimeric cytokine receptors include homodimers (e.g., PDGF receptor $\alpha\alpha$ and $\beta\beta$ isoforms, erythropoietin receptor, MPL [thrombopoietin receptor], and G-CSF receptor), heterodimers whose subunits each have ligand-binding and effector domains (e.g., PDGF receptor $\alpha\beta$ isoform), and multimers having component subunits with disparate functions (e.g., IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, and GM-CSF receptors). Some receptor subunits are common to a plurality of receptors. For example, the AIC2B subunit, which cannot bind ligand on its own but includes an intracellular signal

transduction domain, is a component of IL-3 and GM-CSF receptors. Many cytokine receptors can be placed into one of four related families on the basis of their structures (see Figure) and functions. Hematopoietic receptors, for example, are characterized by the presence of a domain containing conserved cysteine residues and the WSXWS motif (SEQ ID NO:5). Additional domains, including protein kinase domains; fibronectin type III domains; and immunoglobulin domains, which are characterized by disulfide-bonded loops, are present in certain hematopoietic receptors. Cytokine receptor structure has been reviewed by Urdal, Ann. Reports Med. Chem. 26:221-228, 1991 and Cosman, Cytokine 5:95-106, 1993. It is generally believed that under selective pressure for organisms to acquire new biological functions, new receptor family members arose from duplication of existing receptor genes leading to the existence of multi-gene families. Family members thus contain vestiges of the ancestral gene, and these characteristic features can be exploited in the isolation and identification of additional family members. The cytokine receptor superfamily is subdivided as shown in Table 1.

25	<u>Table 1</u>
	<u>Cytokine Receptor Superfamily</u>
	Immunoglobulin family
	CSF-1 receptor
	MGF receptor
30	IL-1 receptor
	PDGF receptor
	Hematopoietin family
	erythropoietin receptor
	G-CSF receptor
35	IL-2 receptor β -subunit
	IL-3 receptor

Table 1. continued

	IL-4 receptor
	IL-5 receptor
	IL-6 receptor
5	IL-7 receptor
	IL-9 receptor
	GM-CSF receptor α -subunit
	GM-CSF receptor β -subunit
	Prolactin receptor
10	CNTF receptor
	Oncostatin M receptor
	Leukemia inhibitory factor receptor
	Growth hormone receptor
	MPL
15	Leptin receptor
	TNF receptor family
	TNF (p80) receptor
	TNF (p60) receptor
	TNFR-RP
20	CD27
	CD30
	CD40
	4-1BB
	OX-40
25	Fas
	NGF receptor
	Other
	IL-2 receptor α -subunit
	IL-15 receptor α -subunit
30	IFN- γ receptor

Cell-surface cytokine receptors are further characterized by the presence of additional domains. These receptors are anchored in the cell membrane by a transmembrane domain characterized by a sequence of hydrophobic amino acid residues (typically about 21-25 residues), which is commonly flanked by positively charged

residues (Lys or Arg). On the opposite end of the protein from the extracellular domain and separated from it by the transmembrane domain is an intracellular domain.

The novel receptor of the present invention was initially identified by the presence of the conserved WSXWS motif (SEQ ID NO:5). Analysis of a human cDNA clone encoding ZCyt2 (SEQ ID NO:1) revealed an open reading frame encoding 380 amino acids (SEQ ID NO:2) comprising an extracellular ligand-binding domain of approximately 315 amino acid residues (residues 25-339 of SEQ ID NO:2), a transmembrane domain of approximately 24 amino acid residues (residues 340-363 of SEQ ID NO:2), and a short intracellular domain of approximately 17 amino acid residues (residues 364-380 of SEQ ID NO:2). Those skilled in the art will recognize that these domain boundaries are approximate and are based on alignments with known proteins and predictions of protein folding. Deletion of residues from the ends of the domains is possible. For example, the core ligand binding region is believed to reside within residues 141-337 of SEQ ID NO:2. Structural analysis indicates that the polypeptide regions from Cys145 through Cys155 and from Cys184 through Cys197 of SEQ ID NO:2 are cysteine loops that are important ligand-binding sites. Relatively small, ligand-binding receptor polypeptides are thus provided by the present invention.

The deduced amino acid sequence of Zcytor2 indicates that it belongs to the same subfamily as the IL-3, IL-5 and GM-CSF receptor α subunits. These α receptor subunits are ligand-specific proteins that combine with a common signalling subunit (β -subunit) to form a signalling complex in the presence of the cognate ligand. The β -subunit for this receptor subfamily has been previously identified in mouse (Itoh et al., Science 247:324-327, 1989; Gorman et al., Proc. Natl. Acad. Sci. USA 87:5459-5463, 1990) and human (Hayashida, et al., Proc. Natl. Acad. Sci. USA 87:9655-9659, 1990). The mouse β -subunit occurs in two isoforms, denoted AIC2A and AIC2B, whereas in human

only one form (denoted β_C) has been identified. β_C is also a member of the hematopoietin receptor family in that it contains a WSXWS motif (SEQ ID NO:5) and a single transmembrane domain. β_C also contains a sizable intracellular domain capable of interacting with cytoplasmic proteins for signal propagation. In the alternative, Zcytor2 may combine with one or more of gp130 (Hibi et al., Cell 63:1149-1157, 1990), the IL-4 α -subunit (Idzerda, et al., J. Exp. Med. 171:861, 1990), or the IL-13 α -subunit (Hilton et al., Proc. Natl. Acad. Sci. USA 93:497-501, 1996) in a tissue specific manner to form dimeric or trimeric complexes. Binding data for Zcytor2 suggest that this receptor subunit may form an IL-13 receptor complex in testes and pituitary that is different from the immune system IL-13 receptor.

Within preferred embodiments of the invention the isolated polynucleotides will hybridize to similar sized regions of SEQ ID NO:1, SEQ ID NO:3, or SEQ ID NO:6, or a sequence complementary thereto, under stringent conditions. In general, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typical stringent conditions are those in which the salt concentration is at least about 0.02 M at pH 7 and the temperature is at least about 60°C. As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for isolating DNA and RNA are well known in the art. It is generally preferred to isolate RNA from testis, including whole testis tissue extracts or testicular cells, such as Sertoli cells, Leydig cells, spermatogonia, or epididymis, although DNA can also be prepared using RNA from other tissues or isolated as genomic DNA. Total RNA can be prepared using guanidine HCl extraction followed by isolation by

centrifugation in a CsCl gradient (Chirgwin et al., Biochemistry 18:52-94, 1979). Poly (A)⁺ RNA is prepared from total RNA using the method of Aviv and Leder (Proc. Natl. Acad. Sci. USA 69:1408-1412, 1972). Complementary DNA (cDNA) is prepared from poly(A)⁺ RNA using known methods. Polynucleotides encoding Zcytor2 polypeptides are then identified and isolated by, for example, hybridization or PCR.

Those skilled in the art will recognize that the sequences disclosed in SEQ ID NOS:1, 2, 6, and 7 represent single alleles of the human and macaque ZCyt2 receptors, respectively. Allelic variants of these sequences can be cloned by probing cDNA or genomic libraries from different individuals according to standard procedures. DNA and protein sequences from an additional human clone are shown in SEQ ID NOS: 3 and 4.

The present invention further provides counterpart receptors and polynucleotides from other species ("species orthologs"). Of particular interest are ZCyt2 receptors from other mammalian species, including murine, porcine, ovine, bovine, canine, feline, equine, and other primate receptors. Species orthologs of the human and macaque ZCyt2 receptors can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA obtained from a tissue or cell type that expresses the receptor. Suitable sources of mRNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from mRNA of a positive tissue or cell line. A receptor-encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial human or macaque cDNA or with one or more sets of degenerate probes based on the disclosed sequences. A cDNA can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent No. 4,683,202).

using primers designed from the sequences disclosed herein. Within an additional method, the cDNA library can be used to transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to the receptor. Similar techniques can also be applied to the isolation of genomic clones.

The present invention also provides isolated receptor polypeptides that are substantially homologous to the receptor polypeptides of SEQ ID NO: 2 or SEQ ID NO:7 and their species orthologs. By "isolated" is meant a protein or polypeptide that is found in a condition other than its native environment, such as apart from blood and animal tissue. In a preferred form, the isolated polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, i.e. greater than 95% pure, more preferably greater than 99% pure. The term "substantially homologous" is used herein to denote polypeptides having 50%, preferably 60%, more preferably at least 80%, sequence identity to the sequences shown in SEQ ID NO:2, 4, or 7 or their species orthologs. Such polypeptides will more preferably be at least 90% identical, and most preferably 95% or more identical to SEQ ID NO:2, 4 or 7 or their species orthologs. Percent sequence identity is determined by conventional methods. See, for example, Altschul et al., Bull. Math. Bio. 48: 603-616, 1986 and Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915-10919, 1992. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "blosum 62" scoring matrix of Henikoff and Henikoff (ibid.) as shown in Table 2 (amino acids are indicated by the standard one-letter codes). The percent identity is then calculated as:

$$\frac{\text{Total number of identical matches}}{[\text{length of the longer sequence plus the}]} \times 100$$

number of gaps introduced into the longer
sequence in order to align the two
sequences]

Sequence identity of polynucleotide molecules is determined by similar methods using a ratio as disclosed above.

Substantially homologous proteins and polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor nature, that is conservative amino acid substitutions (see Table 3) and other substitutions that do not significantly affect the folding or activity of the protein or polypeptide; small deletions, typically of one to about 30 amino acids; and small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, a small linker peptide of up to about 20-25 residues, or a small extension that facilitates purification (an affinity tag), such as a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al., Methods Enzymol. 198:3, 1991), glutathione S transferase (Smith and Johnson, Gene 67:31, 1988), or other antigenic epitope or binding domain. See, in general Ford et al., Protein Expression and Purification 2: 95-107, 1991, which is incorporated herein by reference. DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ).

Table 3

Conservative amino acid substitutions

30	Basic:	arginine
		lysine
		histidine
	Acidic:	glutamic acid
		aspartic acid
35	Polar:	glutamine
		asparagine
	Hydrophobic:	leucine
		isoleucine
		valine

Table 3. continued

	Aromatic:	phenylalanine
		tryptophan
		tyrosine
5	Small:	glycine
		alanine
		serine
		threonine
		methionine

10

Essential amino acids in the receptor polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244, 1081-1085, 1989; Bass et al., Proc. Natl. Acad. Sci. USA 88:4498-4502, 1991). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity (e.g., ligand binding and signal transduction) to identify amino acid residues that are critical to the activity of the molecule. Sites of ligand-receptor interaction can also be determined by analysis of crystal structure as determined by such techniques as nuclear magnetic resonance, crystallography or photoaffinity labeling. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992. The identities of essential amino acids can also be inferred from analysis of homologies with related receptors.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53-57, 1988) or Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-2156, 1989). Briefly, these authors disclose methods for simultaneously randomizing

two or more positions in a polypeptide, selecting for functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988)

10 Mutagenesis methods as disclosed above can be combined with high-throughput screening methods to detect activity of cloned, mutagenized receptors in host cells. Preferred assays in this regard include cell proliferation assays and biosensor-based ligand-binding assays, which
15 are described below. Mutagenized DNA molecules that encode active receptors or portions thereof (e.g., ligand-binding fragments) can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of
20 individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in the art can prepare a variety of
25 polypeptides that are substantially homologous to residues 141 to 337 of SEQ ID NO:2 or allelic variants thereof and retain the ligand-binding properties of the wild-type receptor. Such polypeptides may include additional amino acids from an extracellular ligand-binding domain of a
30 Zcytor2 receptor as well as part or all of the transmembrane and intracellular domains. Such polypeptides may also include additional polypeptide segments as generally disclosed above.

The receptor polypeptides of the present
35 invention, including full-length receptors, receptor fragments (e.g. ligand-binding fragments), and fusion polypeptides can be produced in genetically engineered

host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells. Eukaryotic cells, particularly cultured cells of multicellular organisms, are preferred. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al., *ibid.*, which are incorporated herein by reference.

In general, a DNA sequence encoding a ZCyt2 receptor polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

To direct a ZCyt2 receptor polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) is provided in the expression vector. The secretory signal sequence may be that of the receptor, or may be derived from another secreted protein (e.g., t-PA) or synthesized *de novo*. The secretory signal sequence is joined to the ZCyt2 DNA sequence in the correct reading frame. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the polypeptide

of interest, although certain signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et al., U.S. Patent No. 5,143,830).

5 Cultured mammalian cells are preferred hosts within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection (Wigler et al., Cell 14:725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7:603, 1981; Graham and Van der Eb, Virology 52:456, 10 1973), electroporation (Neumann et al., EMBO J. 1:841-845, 1982), DEAE-dextran mediated transfection (Ausubel et al., eds., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987), and liposome-mediated 15 transfection (Hawley-Nelson et al., Focus 15:73, 1993; Ciccarone et al., Focus 15:80, 1993), which are incorporated herein by reference. The production of recombinant polypeptides in cultured mammalian cells is disclosed, for example, by Levinson et al., U.S. Patent 20 No. 4,713,339; Hagen et al., U.S. Patent No. 4,784,950; Palmiter et al., U.S. Patent No. 4,579,821; and Ringold, U.S. Patent No. 4,656,134, which are incorporated herein by reference. Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), 25 BHK (ATCC No. CRL 1632), BHK 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 1573; Graham et al., J. Gen. Virol. 36:59-72, 1977) and Chinese hamster ovary (e.g. CHO-K1; ATCC No. CCL 61) cell lines. Additional suitable cell lines are known in the art and available from public depositories 30 such as the American Type Culture Collection, Rockville, Maryland. In general, strong transcription promoters are preferred, such as promoters from SV-40 or cytomegalovirus. See, e.g., U.S. Patent No. 4,956,288. Other suitable promoters include those from 35 metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978, which are incorporated herein by reference) and the adenovirus major late promoter.

Drug selection is generally used to select for cultured mammalian cells into which foreign DNA has been inserted. Such cells are commonly referred to as "transfectants". Cells that have been cultured in the presence of the selective agent and are able to pass the gene of interest to their progeny are referred to as "stable transfectants." A preferred selectable marker is a gene encoding resistance to the antibiotic neomycin. Selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems may also be used to increase the expression level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. A preferred amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug resistance genes (e.g. hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can also be used.

Other higher eukaryotic cells can also be used as hosts, including insect cells, plant cells and avian cells. Transformation of insect cells and production of foreign polypeptides therein is disclosed by Guarino et al., U.S. Patent No. 5,162,222; Bang et al., U.S. Patent No. 4,775,624; and WIPO publication WO 94/06463, which are incorporated herein by reference. The use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells has been reviewed by Sinkar et al., J. Biosci. (Bangalore) 11:47-58, 1987.

Fungal cells, including yeast cells, and particularly cells of the genus *Saccharomyces*, can also be used within the present invention, such as for producing receptor fragments or polypeptide fusions. Methods for transforming yeast cells with exogenous DNA and producing

recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311; Kawasaki et al., U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch et al., U.S. Patent No. 5,037,743; and
5 Murray et al., U.S. Patent No. 4,845,075, which are incorporated herein by reference. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (e.g., leucine). A
10 preferred vector system for use in yeast is the POT1 vector system disclosed by Kawasaki et al. (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those
15 from glycolytic enzyme genes (see, e.g., Kawasaki, U.S. Patent No. 4,599,311; Kingsman et al., U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092, which are incorporated herein by reference) and alcohol dehydrogenase genes. See also U.S. Patents Nos.
20 4,990,446; 5,063,154; 5,139,936 and 4,661,454, which are incorporated herein by reference. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*,
25 *Pichia methanolica*, *Pichia guilliermondii* and *Candida maltosa* are known in the art. See, for example, Gleeson et al., J. Gen. Microbiol. 132:3459-3465, 1986 and Cregg, U.S. Patent No. 4,882,279. *Aspergillus* cells may be utilized according to the methods of McKnight et al., U.S.
30 Patent No. 4,935,349, which is incorporated herein by reference. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino et al., U.S. Patent No. 5,162,228, which is incorporated herein by reference. Methods for transforming *Neurospora* are disclosed by
35 Lambowitz, U.S. Patent No. 4,486,533, which is incorporated herein by reference.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of
5 suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required. The growth medium
10 will generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell.

15 Within one aspect of the present invention, a novel receptor is produced by a cultured cell, and the cell is used to screen for ligands for the receptor, including the natural ligand, as well as agonists and antagonists of the natural ligand. To summarize this
20 approach, a cDNA or gene encoding the receptor is combined with other genetic elements required for its expression (e.g., a transcription promoter), and the resulting expression vector is inserted into a host cell. Cells that express the DNA and produce functional receptor are
25 selected and used within a variety of screening systems.

Mammalian cells suitable for use in expressing ZCyt2 receptors and transducing a receptor-mediated signal include cells that express a β -subunit, such as the human β_c subunit. In this regard it is generally
30 preferred to employ a cell that is responsive to other cytokines that bind to receptors in the same subfamily, such as IL-3 or GM-CSF, because such cells will contain the requisite signal transduction pathway(s). It is also preferred to use a cell from the same species as the
35 receptor to be expressed. Within a preferred embodiment, the cell is dependent upon an exogenously supplied hematopoietic growth factor for its proliferation.

Preferred cell lines of this type are the human TF-1 cell line (ATCC number CRL-2003) and the AML-193 cell line (ATCC number CRL-9589), which are GM-CSF-dependent human leukemic cell lines. In the alternative, suitable host
5 cells can be engineered to produce a β -subunit (e.g., β_C) or other cellular component needed for the desired cellular response. For example, the murine cell line BaF3 (Palacios and Steinmetz, Cell 41: 727-734, 1985; Mathey-Prevot et al., Mol. Cell. Biol. 6: 4133-4135, 1986) or a
10 baby hamster kidney (BHK) cell line can be transfected to express the human β_C subunit (also known as KH97) as well as a ZCyt2 receptor. The latter approach is advantageous because cell lines can be engineered to express receptor subunits from any species, thereby
15 overcoming potential limitations arising from species specificity. In the alternative, species orthologs of the human receptor cDNA can be cloned and used within cell lines from the same species, such as a mouse cDNA in the BaF3 cell line. Cell lines that are dependent upon one
20 hematopoietic growth factor, such as GM-CSF, can thus be engineered to become dependent upon a Zcytor2 ligand.

Cells expressing functional receptor are used within screening assays. A variety of suitable assays are known in the art. These assays are based on the detection
25 of a biological response in a target cell. One such assay is a cell proliferation assay. Cells are cultured in the presence or absence of a test compound, and cell proliferation is detected by, for example, measuring incorporation of tritiated thymidine or by colorimetric
30 assay based on the metabolic breakdown of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) (Mosman, J. Immunol. Meth. 65: 55-63, 1983). An alternative assay format uses cells that are further engineered to express a reporter gene. The reporter gene
35 is linked to a promoter element that is responsive to the receptor-linked pathway, and the assay detects activation of transcription of the reporter gene. A preferred

promoter element in this regard is a serum response element, or SRE (see, e.g., Shaw et al., Cell 56:563-572, 1989). A preferred such reporter gene is a luciferase gene (de Wet et al., Mol. Cell. Biol. 7:725, 1987).

5 Expression of the luciferase gene is detected by luminescence using methods known in the art (e.g., Baumgartner et al., J. Biol. Chem. 269:29094-29101, 1994; Schenborn and Goiffin, Promega Notes 41:11, 1993). Luciferase activity assay kits are commercially available

10 from, for example, Promega Corp., Madison, WI. Target cell lines of this type can be used to screen libraries of chemicals, cell-conditioned culture media, fungal broths, soil samples, water samples, and the like. For example, a bank of cell-conditioned media samples can be assayed on a

15 target cell to identify cells that produce ligand. Positive cells are then used to produce a cDNA library in a mammalian expression vector, which is divided into pools, transfected into host cells, and expressed. Media samples from the transfected cells are then assayed, with

20 subsequent division of pools, re-transfection, subculturing, and re-assay of positive cells to isolate a cloned cDNA encoding the ligand.

A natural ligand for the ZCyt2 receptor can also be identified by mutagenizing a cell line expressing

25 the receptor and culturing it under conditions that select for autocrine growth. See WIPO publication WO 95/21930. Within a typical procedure, BaF3 cells expressing ZCyt2 and human β_c are mutagenized, such as with 2-ethylmethanesulfonate (EMS). The cells are then allowed

30 to recover in the presence of IL-3, then transferred to a culture medium lacking IL-3 and IL-4. Surviving cells are screened for the production of a ZCyt2 ligand, such as by adding soluble receptor to the culture medium or by assaying conditioned media on wild-type BaF3 cells and

35 BaF3 cells expressing the receptor.

An additional screening approach provided by the present invention includes the use of hybrid receptor

polypeptides. These hybrid polypeptides fall into two general classes. Within the first class, the intracellular domain of Z-Cytor2, comprising approximately residues 364 to 380 of SEQ ID NO:2, is joined to the ligand-binding domain of a second receptor. It is preferred that the second receptor be a hematopoietic cytokine receptor, such as mpl receptor (Souyri et al., Cell 63: 1137-1147, 1990). The hybrid receptor will further comprise a transmembrane domain, which may be derived from either receptor. A DNA construct encoding the hybrid receptor is then inserted into a host cell. Cells expressing the hybrid receptor are cultured in the presence of a ligand for the binding domain and assayed for a response. This system provides a means for analyzing signal transduction mediated by ZCytor2 while using readily available ligands. This system can also be used to determine if particular cell lines are capable of responding to signals transduced by ZCytor2. A second class of hybrid receptor polypeptides comprise the extracellular (ligand-binding) domain of ZCytor2 (approximately residues 25 to 337 of SEQ ID NO:2) with an intracellular domain of a second receptor, preferably a hematopoietic cytokine receptor, and a transmembrane domain. Hybrid receptors of this second class are expressed in cells known to be capable of responding to signals transduced by the second receptor. Together, these two classes of hybrid receptors enable the use of a broad spectrum of cell types within receptor-based assay systems.

Cells found to express the ligand are then used to prepare a cDNA library from which the ligand-encoding cDNA can be isolated as disclosed above. The present invention thus provides, in addition to novel receptor polypeptides, methods for cloning polypeptide ligands for the receptors.

The tissue specificity of ZCytor2 expression suggests a role in spermatogenesis, a process that is

remarkably similar to the development of blood cells (hematopoiesis). Briefly, spermatogonia undergo a maturation process similar to the differentiation of hematopoietic stem cells. In both systems, the c-kit ligand is involved in the early stages of differentiation. In view of the tissue specificity observed for this receptor, agonists (including the natural ligand) and antagonists have enormous potential in both *in vitro* and *in vivo* applications. Compounds identified as receptor agonists are useful for stimulating proliferation and development of target cells *in vitro* and *in vivo*. For example, agonist compounds are useful as components of defined cell culture media, and may be used alone or in combination with other cytokines and hormones to replace serum that is commonly used in cell culture. Agonists are thus useful in specifically promoting the growth and/or development of testis-derived cells in culture. Agonists and antagonists may also prove useful in the study of spermatogenesis and infertility. Antagonists are useful as research reagents for characterizing sites of ligand-receptor interaction. *In vivo*, receptor agonists may find application in the treatment of male infertility. Antagonists of receptor function may be useful as male contraceptive agents.

Zcytor2 receptor antagonists and ligand-binding polypeptides may also be used to modulate immune functions by blocking the action of IL-13. Of particular interest in this regard is the limiting of unwanted immune responses, such as allergies and asthma. Local administration is preferred to avoid systemic immune suppression. Examples of local administration include topical application to the skin and inhalation. Suitable methods of formulation are known in the art.

Zcytor2 may also be used within diagnostic systems for the detection of circulating levels of ligand. Within a related embodiment, antibodies or other agents that specifically bind to Zcytor2 can be used to detect

circulating receptor polypeptides. Elevated or depressed levels of ligand or receptor polypeptides may be indicative of pathological conditions, including cancer.

5 ZCyt2 receptor polypeptides can be prepared by expressing a truncated DNA encoding residues 141 through 337 of a human Zcytor2 receptor (SEQ ID NO:2 or SEQ ID NO:4) or the corresponding region of a non-human receptor. Additional residues of the receptor may also be included, in particular amino-terminal residues between the
10 predicted mature N-terminus (residue 25 of SEQ ID NO:2 or SEQ ID NO:4) and residue 141, and short C-terminal extensions. It is preferred that the extracellular domain polypeptides be prepared in a form substantially free of transmembrane and intracellular polypeptide segments. For
15 example, the C-terminus of the receptor polypeptide may be at residue 338 or 339 of SEQ ID NO:2 or the corresponding region of an allelic variant or a non-human receptor. A preferred such polypeptide consists of residues 25 to 337 of SEQ ID NO:4. To direct the export of the receptor
20 domain from the host cell, the receptor DNA is linked to a second DNA segment encoding a secretory peptide, such as a t-PA secretory peptide. To facilitate purification of the secreted receptor domain, a C-terminal extension, such as a poly-histidine tag, substance P, FlagTM peptide (Hopp et
25 al., Biotechnology 6:1204-1210, 1988; available from Eastman Kodak Co., New Haven, CT) or another polypeptide or protein for which an antibody or other specific binding agent is available, can be fused to the receptor polypeptide.

30 In an alternative approach, a receptor extracellular domain can be expressed as a fusion with immunoglobulin heavy chain constant regions, typically an F_C fragment, which contains two constant region domains and a hinge region but lacks the variable region. Such
35 fusions are typically secreted as multimeric molecules wherein the Fc portions are disulfide bonded to each other and two receptor polypeptides are arrayed in closed

proximity to each other. Fusions of this type can be used to affinity purify the cognate ligand from solution, as an *in vitro* assay tool, to block signals *in vitro* by specifically titrating out ligand, and as antagonists *in vivo* by administering them parenterally to bind circulating ligand and clear it from the circulation. To purify ligand, a Zcytor2-Ig chimera is added to a sample containing the ligand (e.g., cell-conditioned culture media or tissue extracts) under conditions that facilitate receptor-ligand binding (typically near-physiological temperature, pH, and ionic strength). The chimera-ligand complex is then separated by the mixture using protein A, which is immobilized on a solid support (e.g., insoluble resin beads). The ligand is then eluted using conventional chemical techniques, such as with a salt or pH gradient. In the alternative, the chimera itself can be bound to a solid support, with binding and elution carried out as above. The chimeras may be used *in vivo* to induce infertility. Chimeras with high binding affinity are administered parenterally (e.g., by intramuscular, subcutaneous or intravenous injection). Circulating molecules bind ligand and are cleared from circulation by normal physiological processes. For use in assays, the chimeras are bound to a support via the F_C region and used in an ELISA format.

A preferred assay system employing a ligand-binding receptor fragment uses a commercially available biosensor instrument (BIAcore™, Pharmacia Biosensor, Piscataway, NJ), wherein the receptor fragment is immobilized onto the surface of a receptor chip. Use of this instrument is disclosed by Karlsson, J. Immunol. Methods 145:229-240, 1991 and Cunningham and Wells, J. Mol. Biol. 234:554-563, 1993. A receptor fragment is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within the flow cell. A test sample is passed through the cell. If ligand is present in the sample, it will bind to the

immobilized receptor polypeptide, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film. This system allows the determination of on- and off-rates, from which binding affinity can be calculated, and assessment of stoichiometry of binding.

Ligand-binding receptor polypeptides can also be used within other assay systems known in the art. Such systems include Scatchard analysis for determination of binding affinity (see, Scatchard, Ann. NY Acad. Sci. 51: 660-672, 1949) and calorimetric assays (Cunningham et al., Science 253:545-548, 1991; Cunningham et al., Science 254:821-825, 1991).

A receptor ligand-binding polypeptide can also be used for purification of ligand. The receptor polypeptide is immobilized on a solid support, such as beads of agarose, cross-linked agarose, glass, cellulosic resins, silica-based resins, polystyrene, cross-linked polyacrylamide, or like materials that are stable under the conditions of use. Methods for linking polypeptides to solid supports are known in the art, and include amine chemistry, cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, and hydrazide activation. The resulting media will generally be configured in the form of a column, and fluids containing ligand are passed through the column one or more times to allow ligand to bind to the receptor polypeptide. The ligand is then eluted using changes in salt concentration or pH to disrupt ligand-receptor binding.

Zcytor2 polypeptides can also be used to prepare antibodies that specifically bind to Zcytor2 polypeptides. As used herein, the term "antibodies" includes polyclonal antibodies, monoclonal antibodies, antigen-binding fragments thereof such as F(ab')₂ and Fab fragments, and the like, including genetically engineered antibodies. Antibodies are defined to be specifically binding if they

bind to a Zcytor2 polypeptide with a K_a of greater than or equal to $10^7/M$. The affinity of a monoclonal antibody can be readily determined by one of ordinary skill in the art (see, for example, Scatchard, *ibid.*).

5 Methods for preparing polyclonal and monoclonal antibodies are well known in the art (see for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY, 1989; and Hurrell, J. G. R., Ed., Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC Press, Inc., Boca Raton, FL, 1982, 10 which are incorporated herein by reference). As would be evident to one of ordinary skill in the art, polyclonal antibodies can be generated from a variety of warm-blooded animals such as horses, cows, goats, sheep, dogs, 15 chickens, rabbits, mice, and rats. The immunogenicity of a Zcytor2 polypeptide may be increased through the use of an adjuvant such as Freund's complete or incomplete adjuvant. A variety of assays known to those skilled in the art can be utilized to detect antibodies which 20 specifically bind to Zcytor2 polypeptides. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988. Representative examples of such assays include: concurrent immunoelectrophoresis, radio- 25 immunoassays, radio-immunoprecipitations, enzyme-linked immunosorbent assays (ELISA), dot blot assays, inhibition or competition assays, and sandwich assays.

 Antibodies to Zcytor2 are may be used for tagging cells that express the receptor, for affinity 30 purification, within diagnostic assays for determining circulating levels of soluble receptor polypeptides, and as antagonists to block ligand binding and signal transduction *in vitro* and *in vivo*.

 The invention is further illustrated by the 35 following non-limiting examples.

Example 1

A cDNA library was prepared from human placental poly A⁺ RNA provided as a control in a MarathonTM cDNA Amplification Kit (Clontech, Palo Alto, CA) using the
5 protocol provided by the manufacturer. This cDNA was used as template in polymerase chain reactions to generate DNA encoding human Zcytor2.

Primers were designed from the sequences of two expressed sequence tags (ESTs) in a DNA sequence database.
10 Analysis of the EST sequences suggested that they represented the 5' and 3' ends of a cDNA encoding a cytokine receptor. One pair of primers, designated ZG9801 (SEQ ID NO:8) and ZG9941 (SEQ ID NO:9), were designed to be used in a 5' RACE (rapid amplification of cDNA ends)
15 reaction. A second pair, designated ZG9803 (SEQ ID NO:10) and ZG9937 (SEQ ID NO:11), were designed to be used in a 3' RACE reaction. A third pair of primers, designated ZG9800 (SEQ ID NO:12) and ZG9802 (SEQ ID NO:13), were designed to amplify the region spanning the two ESTs. A
20 fourth pair of primers, AP1 (SEQ ID NO:14) and AP2 (SEQ ID NO:15), were supplied with the amplification kit or synthesized.

PCR amplification was carried out according to the instruction manual supplied with the kit, with certain
25 modifications to the protocol. For the 5' and 3' RACE reactions, fifty pmol of each primer was used in each reaction. Each cDNA template was initially amplified using the appropriate gene-specific primer (ZG9801 or ZG9803) for 10 cycles. Primer AP1 was then added, and the
30 reaction was continued for 25 cycles. The reaction mixture was incubated in a Hybaid OmniGene Temperature Cycling System (National Labnet Co., Woodbridge, NY) for 1 minute at 95°C, then for 10 cycles of 60°C, 30 seconds; 72°C, 2 minutes; 95°C, 30 seconds. The mixture was held at 60
35 °C, and 50 pmol of primer AP1 was added, and the reaction was continued for 25 cycles of 60°C, 30 seconds; 72°C, 2 minutes; 95°C, 30 seconds; followed by a 7 minute

incubation at 72°C. The internal fragment was amplified under the same conditions using gene-specific primers (9800 and 9802), but AP1 was omitted. Reaction products were analyzed by electrophoresis on a 1% agarose gel. A
5 discreet band was obtained for the internal fragment. The 5' and 3' RACE products were smears on the gel.

The 5' and 3' RACE products were purified using a PCR purification kit (Qiagen Inc., Chatsworth, CA) and used in nested PCR reactions. Each template was combined
10 with 50 pmol of the appropriate specific primer (ZG9941 or ZG9937) and 50 pmol of primer AP2. Reactions were run for 30 cycles of 95°C, 1 minute; 60°C, 30 seconds; 72°C, 3.5 minutes; then incubated at 72°C for 7 minutes. The reaction products were analyzed by electrophoresis on a 1%
15 agarose gel. One discreet band was obtained for each reaction.

The 5' and 3' products from the nested PCR reactions and the internal fragment from the initial Marathon™ PCR reaction were gel purified using a Qiagen
20 Gel Extraction Kit.

The internal fragment was subcloned using a Stratagene (La Jolla, CA) PCR-Script™ SK(+) Cloning Kit according to the manufacturer's instructions, with 10 µl H₂O added to each reaction. The ligated DNA was then
25 purified using CENTRI-SEP columns (Princeton Separations, Adelphia, NJ) to increase the efficiency of transformation. The resulting vector was used to transform *E. coli* Electromax DH10B™ cells (Gibco BRL, Gaithersburg, MD) by electroporation.

30 Colonies were screened by PCR using gene-specific primers. Individual white colonies representing recombinants were picked and added to microcentrifuge tubes by swirling the toothpick with the colony on it in a tube containing 19.5 µl H₂O, 2.5 µl 10x Taq polymerase
35 buffer (Boehringer Mannheim, Indianapolis, IN), 0.5 µl 10 mM dNTPs, 1.0 µl ZG9800 (SEQ ID NO:12) (20 pmol/µl), 10 µl ZG9802 (SEQ ID NO:13) (20 pmol/µl), and 0.5 µl Taq

polymerase. Cells were streaked out on a master plate to use for starting cultures. Amplification reactions were incubated at 96°C for 45 seconds to lyse the bacteria and expose the plasmid DNA, then run for 25 cycles of 96°C, 45 seconds; 55°C, 45 seconds; 72°C, 2 minutes to amplify cloned inserts. Products were analyzed by electrophoresis on a 1% agarose gel. One clone was identified as positive, and a plasmid template was prepared for sequencing using a QIAwell™ 8 Plasmid Kit (Qiagen Inc.).

The 5' RACE product, the 3' RACE product, the internal fragment and the internal fragment subclone were sequenced on an Applied Biosystems™ model 373 DNA sequencer (Perkin-Elmer Corporation, Norwalk, CT) using either an AmpliTaq® DyeDeoxy™ Terminator Cycle Sequencing Kit (Perkin-Elmer Corp.) or an ABI PRISM™ Dye Terminator Cycle Sequencing Core Kit (Perkin-Elmer Corp.). Oligonucleotides used in the PCR reactions were used as sequencing primers. In addition, primers ZG9850 (SEQ ID NO:16), ZG9851 (SEQ ID NO:17), ZG9852 (SEQ ID NO:18) and ZG9919 (SEQ ID NO:19) were used. Sequencing reactions were carried out in a Hybaid OmniGene Temperature Cycling System. Sequencher™ 3.0 sequence analysis software (Gene Codes Corporation, Ann Arbor, MI) was used for data analysis. Although the internal fragment subclone contained the entire coding sequence for the receptor, a composite sequence was constructed from all templates to include additional 5' and 3' untranslated sequence from the RACE products that was not present in the internal subclone. The full sequence is disclosed in SEQ ID NO:1.

A human cDNA was isolated by PCR using oligonucleotide primers specific for the gene sequence and containing restriction sites for subsequent manipulation of the DNA. Specific DNA was amplified from a human testis cDNA library using primers ZG10317 (SEQ ID NO:20) and ZG10319 (SEQ ID NO:21). 10 ng of template DNA was combined with 20 pmol of each primer, 5 µl of 10X buffer (Takara Shuzo Co., Ltd., Otsu, Shiga, Japan), 1 µl of

ExTaq DNA polymerase (Takara Shuzo Co., Ltd.), and 200 μ M dNTPs. The reaction was run for 30 cycles of 95°C, 30 seconds; 55°C, 30 seconds, and 68°C, 2 minutes; then incubated at 68°C for 10 minutes. A fragment of approximately 1200 bp was recovered using a Wizard™ PCR Preps Purification System (Promega Corp., Madison, WI), cleaved with Xho I and Xba I, and a 1200 bp fragment was recovered by precipitation with ethanol.

The 1200 bp fragment was ligated into pHZ200, a vector comprising the mouse metallothionein-1 promoter, the bacteriophage T7 promoter flanked by multiple cloning banks containing unique restriction sites for insertion of coding sequences, the human growth hormone terminator, the bacteriophage T7 terminator, an *E. coli* origin of replication, a bacterial beta lactamase gene, and a mammalian selectable marker expression unit comprising the SV40 promoter and origin, a DHFR gene, and the SV40 transcription terminator. Plasmid pHZ200 was cleaved with Sal I and Xba I and was ligated to the Zcytor2 fragment.

The sequence of the human testis cDNA clone and the deduced amino acid sequence are shown in SEQ ID NO:3 and SEQ ID NO:4, respectively. The deduced amino acid sequence differs from that shown in SEQ ID NO:2 at residues 65, 180, and 259.

25

Example 2

Human Multiple Tissue Northern Blots (Human I, Human II, and Human III from Clontech) were probed to determine the tissue distribution of ZCyt2 expression. A probe was prepared by PCR. Single stranded DNA was prepared from K-562 mRNA (obtained from Clontech) using a RT-PCR kit (Stratagene Cloning Systems, La Jolla, CA) for use as template. 10 ng of template DNA was combined with 20 pmol of each of primers ZG9820 (SEQ ID NO:22) and ZG9806 (SEQ ID NO:23), 5 μ l of 10X buffer (Clontech), 1 μ l of KlenTaq DNA polymerase (Clontech), and 200 μ M dNTPs. The reaction was run for 30 cycles of 95°C, 30 seconds;

55°C, 30 seconds, and 68°C, 2 minutes; then incubated at 68°C for 10 minutes. The resulting DNA was purified by gel electrophoresis and ligated into pGEM®A/T (Promega Corp.). The resulting plasmid was used as a PCR template to generate the probe using the same reaction conditions described above for the K-562 template. DNA was purified by gel electrophoresis and labeled with ³²p by random priming. The blots were prehybridized in ExpressHyb™ hybridization solution (Clontech) at 65°C for 1-6 hours, then hybridized in ExpressHyb™ solution containing 2 x 10⁶ cpm/ml of probe at 65°C for from 1.5 hour to overnight. After hybridization the blots were washed at 50°C in 0.1X SSC, 0.1% SDS. A transcript of approximately 1.5 kb was seen only in testis.

Example 3

A cDNA encoding a soluble human ZCyt2 receptor polypeptide was prepared by PCR. Human cDNA was prepared from a human testis cDNA library. DNA was amplified by PCR using 10 pmol each of oligonucleotide primers ZG10320 (SEQ ID NO:24) and ZG10318 (SEQ ID NO:25). 10 ng of template DNA was combined with 20 pmol of each primer, 5 µl of 10X buffer (Takara Shuzo Co., Ltd.), 1 µl of Taq DNA polymerase (Boehringer Mannheim), and 200 µM dNTPs. The reaction was run for 30 cycles of 95°C, 30 seconds; 55°C, 30 seconds, and 68°C, 2 minutes; then incubated at 68°C for ten minutes. PCR products were separated by electrophoresis on a low melting point agarose gel (Boehringer Mannheim) and purified using a Wizard™ PCR Preps Purification System (Promega Corp.). The fragment was inserted into plasmid HSRT9 that had been cleaved with Bgl II and Xho I. HSRT9 is a mammalian cell expression vector derived from pHZ200 that contains a tissue plasminogen activator (t-PA) secretory signal sequence and a sequence encoding a C-terminal polyhistidine tag downstream of the MT-1 promoter. The resulting construct

encoded a t-PA secretory peptide, human Zcytor2 residues 25--339 (SEQ ID NO:4), and a polyhistidine tag.

5 The soluble receptor expression vector is transfected into BHK 570 cells (ATCC No. CRL-10314) by liposome-mediated transfection (LIPOFECTAMINE™ Reagent, Life Technologies, Gaithersburg, MD). Transfectants are cultured in the presence of methotrexate to select and amplify the transfected DNA. Soluble receptor polypeptide is recovered from conditioned culture media on nickel
10 affinity purification columns (e.g., Talon spin columns from Clontech Laboratories). Columns are washed at neutral pH, and protein is eluted using a decreasing pH gradient or an imidazole gradient. Receptor monomers elute at about pH 6.0-6.3 of 50 mM imidazole, and receptor
15 dimers elute at about pH 5.0-5.3 or 100 mM imidazole. In the alternative, batch purification can be employed.

Example 4

20 A cDNA library was prepared from a non-human primate. Testis tissue was obtained from a 13-year-old Celebus macaque. Total RNA was prepared from the tissue by the CsCl method (Chirgwin et al., Biochemistry 18:52-94, 1979). Poly(A)+ RNA was prepared from the total RNA by oligo(dT) cellulose chromatography (Aviv and Leder,
25 Proc. Natl. Acad. Sci. USA 69:1408-1412, 1972). Double-stranded DNA was prepared from 1 µg of mRNA using a commercially available kit (Clontech Marathon™ cDNA amplification kit).

30 The macaque cDNA was amplified by PCR using a standard adapter-primer and primers derived from the human receptor cDNA sequence. Individual PCR mixtures (50 µl total volume) contained 5 µl template DNA, 5 µl 10X buffer (Clontech), 200 µM dNTPs (Perkin Elmer, CITY), 1 µl each of 10 pmol/µl primer AP1 (Clontech) and one of the
35 primers (20 pmol/µl) shown in Table 4, and 1 µl of Klentaq DNA polymerase (Clontech). The reactions were run for 3 cycles of 94°C, 30 seconds; 65°C, 30 seconds; 68°C, 30

seconds; 3 cycles of 94°C, 30 seconds; 60°C, 30 seconds; 68°C, 30 seconds; 3 cycles of 94°C, 30 seconds; 55°C, 30 seconds; 68°C, 30 seconds; and 30 cycles of 94°C, 30 seconds; 50°, 30 seconds; 68°C, 30 seconds; followed by a
 5 68°C incubation for 10 minutes.

Table 4

	<u>Reaction No.</u>	<u>Primer No.</u>	<u>Primer SEQ ID NO.</u>
10	1	9800	12
	2	9820	22
	3	9941	9
	4	9801	8
	5	9882	26
15	6	10082	27
	7	9850	16
	8	9919	16
	9	10083	28
	10	9803	10
20	11	10081	29
	12	9881	30
	13	9937	11
	14	9806	23
	15	9802	13

25

PCR products were electrophoresed on an agarose gel. The gel was stained with ethidium bromide and viewed under ultraviolet light. Bands from reactions amplified with primers 9800 and 9802 were of the expected size.

30

A second set of PCR reactions was run using the macaque cDNA (1:250 dilution) or first round PCR products from reactions 1, 2, 14 or 15 (Table 4) as templates. the first round PCR products were purified using a Wizard™ PCR Preps Purification System (Promega Corp.) prior to use. 5
 35 µl of template DNA was combined with other components as shown in Table 5. 1 µl of Klentaq DNA polymerase (Clontech) was added to each mixture. Reaction conditions

were as specified above. Reaction products were electrophoresed on an agarose gel, stained with ethidium bromide, and visualized under UV light.

5

Table 5

Rxn. No.	Template	10x Buffer	dNTPs	Primer 1	Primer 2	H ₂ O
1	macaque	5 µl	0.5 µl	--	--	36.5 µl
2	macaque	5 µl	0.5 µl	9800	--	36.5 µl
3	macaque	5 µl	0.5 µl	9802	--	36.5 µl
4	macaque	5 µl	0.5 µl	9800	AP1	36.5 µl
5	macaque	5 µl	0.5 µl	9802	AP1	36.5 µl
6	macaque	5 µl	0.5 µl	AP1	--	36.5 µl
7	macaque	5 µl	0.5 µl	AP1	3'GP3DH	36.5 µl
8	macaque	5 µl	0.5 µl	AP1	5'GP3DH	36.5 µl
9	#14	5 µl	0.5 µl	AP1	9806	36.5 µl
10	#15	5 µl	0.5 µl	AP1	9802	36.5 µl
11	#1	5 µl	0.5 µl	AP1	9800	36.5 µl
12	#2	5 µl	0.5 µl	AP1	9820	36.5 µl

Partial DNA and deduced amino acid sequences of macaque Zcytor2 cDNA are shown in SEQ ID NO:6 and SEQ ID NO:7. Alignment of the human and partial macaque sequences showed an amino acid sequence identity of 92% and a nucleotide sequence identity of 96%.

Example 5

15

An expression vector encoding a human Zcytor2-IgG fusion protein was constructed. The fusion comprised the extracellular domain of Zcytor2 fused at its C-terminus (residue 339 of SEQ ID NO:4) to the hinge region of the Fc portion of an IgG_γ1 (Ellison et al., Nuc. Acids Res. 10:4071-4079, 1982). The hinge region was modified to replace a cysteine residue with serine to avoid unpaired cysteines upon dimerization of the fusion

20

protein. A human t-PA secretory peptide was used to direct secretion of the fusion.

A human Zcytor2 DNA was prepared from a testis cDNA library by PCR using oligonucleotide primers ZG10320 (SEQ ID NO:24) and ZG10389 (SEQ ID NO:31). Twenty pmol of each primer was combined with 1 μ l (10 ng) of template DNA, 10 μ l of 2.5 mM dNTPs (Perkin-Elmer Corp.), 10 μ l of 10X buffer (Klentaq PCR buffer, Clontech), 2 μ l of Klentaq DNA polymerase (Clontech), and 70.8 μ l H₂O. The reaction was run for 35 cycles of 94°C, 1 minute; 55°C, 1 minute; and 72°C, 2 minutes; followed by a 7 minute incubation at 72°C. The reaction products were extracted with phenol/CHCl₃, precipitated with ethanol, and digested with BglII. The DNA was electrophoresed on a agarose gel, and a 941 bp fragment was electrophoretically eluted from a gel slice, purified by phenol/CHCl₃ extraction, and precipitated with ethanol.

A human IgG₁ clone was isolated from a human fetal liver cDNA library (Clontech) by PCR using oligonucleotide primers ZG10314 (SEQ ID NO:32) and ZG10315 (SEQ ID NO:33). The former primer introduced a BglII site into the hinge region (changing the third residue of the hinge region from Lys to Arg) and replaced the fifth residue of the hinge region (Cys) with Ser. PCR was carried out essentially as described above for the Zcytor2 extracellular domain sequence. The DNA was digested with EcoRI and XbaI, and a 0.7 kb fragment was recovered by agarose gel electrophoresis, electroelution, phenol/CHCl₃ extraction, and ethanol precipitation. The IgG-encoding fragment and an XbaI-EcoRI linker were ligated into Zem229R (ATCC Accession No. 69447) that had been digested with EcoRI and treated with calf intestinal phosphatase. The resulting plasmid was digested with BglII and XbaI, and a 950 bp fragment was recovered by agarose gel electrophoresis, electroelution, phenol/CHCl₃ extraction, and ethanol precipitation.

To construct an expression vector for the Zcytor2-IgG fusion, a Zem229R vector containing a human t-PA secretory signal sequence joined to a human thrombopoietin sequence (disclosed in copending, commonly assigned U.S. Patent Application Serial No. 08/347,029) was cleaved with BglII and XbaI. The fragment comprising the vector and t-PA secretory signal sequence was recovered and ligated to the IgG fragment. The Zcytor2 fragment was then ligated into this construct at the BglII site. The resulting plasmid was screened for the desired insert orientation. A plasmid with the desired orientation was designated h-Zcytor-2/IgG #709. Sequence analysis revealed a PCR-generated substitution resulting in an alanine codon instead of a valine codon at position 308 of SEQ ID NO:3.

Plasmid h-Zcytor-2/IgG was transfected into BHK-570 cells by liposome-mediated transfection (LIPOFECTAMINE™ Reagent, Life Technologies, Gaithersburg, MD). Transfectants were cultured in medium containing 1 µM methotrexate for 10 days.

Example 6

The binding of ¹²⁵I-IL-13 to wild-type and Zcytor2-transfected BHK, TF-1, and BaF3 cells was determined. BHK cells were assayed in 6-well culture plates. TF-1 and BaF3 cells were assayed in microcentrifuge tubes. Cells were combined with 500 µl of either solution A (15 ml of binding buffer [RPMI containing 20 mM Tris pH7.4, 0.05% NaN₃, and 3 mg/ml BSA] plus 263 µl of ¹²⁵I-IL-13 [5.7 x 10⁷ cpm/ml]) or solution B (solution A containing 15 µl of cold 25 µg/ml IL-13). After a 2-hour incubation, cells were washed three times with 500 µl binding buffer and lysed in 500 µl of 400 mM NaOH. Lysates were transferred to tubes for gamma counting. BHK cells transfected to express Zcytor2 were found to specifically bind significant amounts of IL-13.

In further experiments, binding of labeled IL-13 was found to be inhibited by IL-13 but not by IL-4.

Saturation binding analysis indicated that Zcytor2 expressed in BHK cells bound ^{125}I -IL-13 with a K_d of 590 ± 359 pM.

To determine if a soluble Zcytor2-IgG fusion could specifically bind IL-13, 1 μg of purified fusion protein was incubated in 200 μl of binding buffer containing 1 nM ^{125}I -IL-13 \pm 100 nM unlabeled IL-13 or IL-4. After two hours at room temperature with mixing, 25 μl of protein A-Sepharose was added, and the mixtures were incubated for an additional hour. The Sepharose was washed three times and collected by centrifugation. Bound ^{125}I -IL-13 was determined by gamma counting. The fusion protein was found to bind significant amounts of labeled IL-13, which was blocked by excess unlabeled IL-13 but not by IL-4.

Binding of labeled IL-13 by BHK/Zcytor2 cells was measured in the presence and absence of the soluble Zcytor2-IgG fusion (0.005 - 5 ng/ml) or unlabeled IL-13. Binding was assayed essentially as described above. Both IL-13 and the fusion protein were found to inhibit binding of labeled IL-13 to the cells.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: ZymoGenetics, Inc.
1201 Eastlake Avenue East
Seattle
WA
USA
98102

(ii) TITLE OF INVENTION: Testis-Specific Receptor

(iii) NUMBER OF SEQUENCES: 33

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: ZymoGenetics, Inc.
(B) STREET: 1201 Eastlake Avenue East
(C) CITY: Seattle
(D) STATE: WA
(E) COUNTRY: USA
(F) ZIP: 98102

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0. Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Parker, Gary E.
(B) REGISTRATION NUMBER: 31,648
(C) REFERENCE/DOCKET NUMBER: 95-33

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206-442-6673
(B) TELEFAX: 206-442-6678

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1289 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 49..1191

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCCCCGCCC GGGAGAGAGG CAATATCAAG GTTTTAAATC TCGGAGAA ATG GCT TTC	57
Met Ala Phe	
1	
GTT TGC TTG GCT ATC GGA TGC TTA TAT ACC TTT CTG ATA AGC ACA ACA	105
Val Cys Leu Ala Ile Gly Cys Leu Tyr Thr Phe Leu Ile Ser Thr Thr	
5 10 15	
TTT GGC TGT ACT TCA TCT TCA GAC ACC GAG ATA AAA GTT AAC CCT CCT	153
Phe Gly Cys Thr Ser Ser Ser Asp Thr Glu Ile Lys Val Asn Pro Pro	
20 25 30 35	
CAG GAT TTT GAG ATA GTG GAT CCC GGA TAC TTA GGT TAT CTC TAT TTG	201
Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr Leu Gly Tyr Leu Tyr Leu	
40 45 50	
CAA TGG CAA CCC CCA CTG TCT CTG GAT CAT TTT AAG GAA TGC ACA GTG	249
Gln Trp Gln Pro Pro Leu Ser Leu Asp His Phe Lys Glu Cys Thr Val	
55 60 65	
GAA TAT GAA CTA AAA TAC CGA AAC ATT GGT AGT GAA ACA TGG AAG ACC	297
Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly Ser Glu Thr Trp Lys Thr	
70 75 80	
ATC ATT ACT AAG AAT CTA CAT TAC AAA GAT GGG TTT GAT CTT AAC AAG	345
Ile Ile Thr Lys Asn Leu His Tyr Lys Asp Gly Phe Asp Leu Asn Lys	
85 90 95	

GGC ATT GAA GCG AAG ATA CAC ACG CTT TTA CCA TGG CAA TGC ACA AAT Gly Ile Glu Ala Lys Ile His Thr Leu Leu Pro Trp Gln Cys Thr Asn 100 105 110 115	393
GGA TCA GAA GTT CAA AGT TCC TGG GCA GAA ACT ACT TAT TGG ATA TCA Gly Ser Glu Val Gln Ser Ser Trp Ala Glu Thr Thr Tyr Trp Ile Ser 120 125 130	441
CCA CAA GGA ATT CCA GAA ACT AAA GTT CAG GAT ATG GAT TGC GTA TAT Pro Gln Gly Ile Pro Glu Thr Lys Val Gln Asp Met Asp Cys Val Tyr 135 140 145	489
TAC AAT TGG CAA TAT TTA CTC TGT TCT TGG AAA CCT GGC ATA GGT GTA Tyr Asn Trp Gln Tyr Leu Leu Cys Ser Trp Lys Pro Gly Ile Gly Val 150 155 160	537
CTT CTT GAT ACC AAT TAC AAC TTG TTT TAC TGG TAT GAG GGC TTG GAT Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr Trp Tyr Glu Gly Leu Asp 165 170 175	585
CAT GCA TTA CAG TGT GTT GAT TAC ATC AAG GCT GAT GGA CAA AAT ATA His Ala Leu Gln Cys Val Asp Tyr Ile Lys Ala Asp Gly Gln Asn Ile 180 185 190 195	633
GGA TGC AGA TTT CCC TAT TTG GAG GCA TCA GAC TAT AAA GAT TTC TAT Gly Cys Arg Phe Pro Tyr Leu Glu Ala Ser Asp Tyr Lys Asp Phe Tyr 200 205 210	681
ATT TGT GTT AAT GGA TCA TCA GAG AAC AAG CCT ATC AGA TCC AGT TAT Ile Cys Val Asn Gly Ser Ser Glu Asn Lys Pro Ile Arg Ser Ser Tyr 215 220 225	729
TTC ACT TTT CAG CTT CAA AAT ATA GTT AAA CCT TTG CCG CCA GTC TAT Phe Thr Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro Pro Val Tyr 230 235 240	777
CTT ACT TTT ACT CGG GAG AGT TCA TGT GAA ATT AAG CTG AAA TGG AGC Leu Thr Phe Thr Arg Glu Ser Ser Cys Glu Ile Lys Leu Lys Trp Ser 245 250 255	825
ATA CCT TTG GGA CCT ATT CCA GCA AGG TGT TTT GAT TAT GAA ATT GAG Ile Pro Leu Gly Pro Ile Pro Ala Arg Cys Phe Asp Tyr Glu Ile Glu 260 265 270 275	873

ATC AGA GAA GAT GAT ACT ACC TTG GTG ACT GCT ACA GTT GAA AAT GAA Ile Arg Glu Asp Asp Thr Thr Leu Val Thr Ala Thr Val Glu Asn Glu 280 285 290	921
ACA TAC ACC TTG AAA ACA ACA AAT GAA ACC CGA CAA TTA TGC TTT GTA Thr Tyr Thr Leu Lys Thr Thr Asn Glu Thr Arg Gln Leu Cys Phe Val 295 300 305	969
GTA AGA AGC AAA GTG AAT ATT TAT TGC TCA GAT GAC GGA ATT TGG AGT Val Arg Ser Lys Val Asn Ile Tyr Cys Ser Asp Asp Gly Ile Trp Ser 310 315 320	1017
GAG TGG AGT GAT AAA CAA TGC TGG GAA GGT GAA GAC CTA TCG AAG AAA Glu Trp Ser Asp Lys Gln Cys Trp Glu Gly Glu Asp Leu Ser Lys Lys 325 330 335	1065
ACT TTG CTA CGT TTC TGG CTA CCA TTT GGT TTC ATC TTA ATA TTA GTT Thr Leu Leu Arg Phe Trp Leu Pro Phe Gly Phe Ile Leu Ile Leu Val 340 345 350 355	1113
ATA TTT GTA ACC GGT CTG CTT TTG CGT AAG CCA AAC ACC TAC CCA AAA Ile Phe Val Thr Gly Leu Leu Leu Arg Lys Pro Asn Thr Tyr Pro Lys 360 365 370	1161
ATG ATT CCA GAA TTT TTC TGT GAT ACA TGAAGACTTT CCATATCAAG Met Ile Pro Glu Phe Phe Cys Asp Thr 375 380	1208
AGACATGGTA TTGACTCAAC AGTTTCCAGT CATGGCCAAA TGTTCATAT GAGTCTCAAT	1268
AAACTGAATT TTTCTTGCGA A	1289

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Phe Val Cys Leu Ala Ile Gly Cys Leu Tyr Thr Phe Leu Ile
 1 5 10 15
 Ser Thr Thr Phe Gly Cys Thr Ser Ser Ser Asp Thr Glu Ile Lys Val
 20 25 30
 Asn Pro Pro Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr Leu Gly Tyr
 35 40 45
 Leu Tyr Leu Gln Trp Gln Pro Pro Leu Ser Leu Asp His Phe Lys Glu
 50 55 60
 Cys Thr Val Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly Ser Glu Thr
 65 70 75 80
 Trp Lys Thr Ile Ile Thr Lys Asn Leu His Tyr Lys Asp Gly Phe Asp
 85 90 95
 Leu Asn Lys Gly Ile Glu Ala Lys Ile His Thr Leu Leu Pro Trp Gln
 100 105 110
 Cys Thr Asn Gly Ser Glu Val Gln Ser Ser Trp Ala Glu Thr Thr Tyr
 115 120 125
 Trp Ile Ser Pro Gln Gly Ile Pro Glu Thr Lys Val Gln Asp Met Asp
 130 135 140
 Cys Val Tyr Tyr Asn Trp Gln Tyr Leu Leu Cys Ser Trp Lys Pro Gly
 145 150 155 160
 Ile Gly Val Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr Trp Tyr Glu
 165 170 175
 Gly Leu Asp His Ala Leu Gln Cys Val Asp Tyr Ile Lys Ala Asp Gly
 180 185 190
 Gln Asn Ile Gly Cys Arg Phe Pro Tyr Leu Glu Ala Ser Asp Tyr Lys
 195 200 205
 Asp Phe Tyr Ile Cys Val Asn Gly Ser Ser Glu Asn Lys Pro Ile Arg
 210 215 220
 Ser Ser Tyr Phe Thr Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro
 225 230 235 240

Pro	Val	Tyr	Leu	Thr	Phe	Thr	Arg	Glu	Ser	Ser	Cys	Glu	Ile	Lys	Leu
				245					250					255	
Lys	Trp	Ser	Ile	Pro	Leu	Gly	Pro	Ile	Pro	Ala	Arg	Cys	Phe	Asp	Tyr
			260					265					270		
Glu	Ile	Glu	Ile	Arg	Glu	Asp	Asp	Thr	Thr	Leu	Val	Thr	Ala	Thr	Val
		275					280					285			
Glu	Asn	Glu	Thr	Tyr	Thr	Leu	Lys	Thr	Thr	Asn	Glu	Thr	Arg	Gln	Leu
	290					295					300				
Cys	Phe	Val	Val	Arg	Ser	Lys	Val	Asn	Ile	Tyr	Cys	Ser	Asp	Asp	Gly
305					310					315					320
Ile	Trp	Ser	Glu	Trp	Ser	Asp	Lys	Gln	Cys	Trp	Glu	Gly	Glu	Asp	Leu
				325					330					335	
Ser	Lys	Lys	Thr	Leu	Leu	Arg	Phe	Trp	Leu	Pro	Phe	Gly	Phe	Ile	Leu
			340					345					350		
Ile	Leu	Val	Ile	Phe	Val	Thr	Gly	Leu	Leu	Leu	Arg	Lys	Pro	Asn	Thr
		355					360					365			
Tyr	Pro	Lys	Met	Ile	Pro	Glu	Phe	Phe	Cys	Asp	Thr				
	370					375					380				

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1167 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
(B) LOCATION: 10..1152

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GATCCGCCC ATG GCT TTC GTT TGC TTG GCT ATC GGA TGC TTA TAT ACC Met Ala Phe Val Cys Leu Ala Ile Gly Cys Leu Tyr Thr 1 5 10	48
TTT CTG ATA AGC ACA ACA TTT GGC TGT ACT TCA TCT TCA GAC ACC GAG Phe Leu Ile Ser Thr Thr Phe Gly Cys Thr Ser Ser Ser Asp Thr Glu 15 20 25	96
ATA AAA GTT AAC CCT CCT CAG GAT TTT GAG ATA GTG GAT CCC GGA TAC Ile Lys Val Asn Pro Pro Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr 30 35 40 45	144
TTA GGT TAT CTC TAT TTG CAA TGG CAA CCC CCA CTG TCT CTG GAT CAT Leu Gly Tyr Leu Tyr Leu Gln Trp Gln Pro Pro Leu Ser Leu Asp His 50 55 60	192
TTT AAG GAA TAC ACA GTG GAA TAT GAA CTA AAA TAC CGA AAC ATT GGT Phe Lys Glu Tyr Thr Val Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly 65 70 75	240
AGT GAA ACA TGG AAG ACC ATC ATT ACT AAG AAT CTA CAT TAC AAA GAT Ser Glu Thr Trp Lys Thr Ile Ile Thr Lys Asn Leu His Tyr Lys Asp 80 85 90	288
GGG TTT GAT CTT AAC AAG GGC ATT GAA GCG AAG ATA CAC ACG CTT TTA Gly Phe Asp Leu Asn Lys Gly Ile Glu Ala Lys Ile His Thr Leu Leu 95 100 105	336
CCA TGG CAA TGC ACA AAT GGA TCA GAA GTT CAA AGT TCC TGG GCA GAA Pro Trp Gln Cys Thr Asn Gly Ser Glu Val Gln Ser Ser Trp Ala Glu 110 115 120 125	384
ACT ACT TAT TGG ATA TCA CCA CAA GGA ATT CCA GAA ACT AAA GTT CAG Thr Thr Tyr Trp Ile Ser Pro Gln Gly Ile Pro Glu Thr Lys Val Gln 130 135 140	432
GAT ATG GAT TGC GTA TAT TAC AAT TGG CAA TAT TTA CTC TGT TCT TGG Asp Met Asp Cys Val Tyr Tyr Asn Trp Gln Tyr Leu Leu Cys Ser Trp 145 150 155	480
AAA CCT GGC ATA GGT GTA CTT CTT GAT ACC AAT TAC AAC TTG TTT TAC Lys Pro Gly Ile Gly Val Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr 160 165 170	528

TGG TAT GAG GGC TTG GAT CTT GCA TTA CAG TGT GTT GAT TAC ATC AAG Trp Tyr Glu Gly Leu Asp Leu Ala Leu Gln Cys Val Asp Tyr Ile Lys 175 180 185	576
GCT GAT GGA CAA AAT ATA GGA TGC AGA TTT CCC TAT TTG GAG GCA TCA Ala Asp Gly Gln Asn Ile Gly Cys Arg Phe Pro Tyr Leu Glu Ala Ser 190 195 200 205	624
GAC TAT AAA GAT TTC TAT ATT TGT GTT AAT GGA TCA TCA GAG AAC AAG Asp Tyr Lys Asp Phe Tyr Ile Cys Val Asn Gly Ser Ser Glu Asn Lys 210 215 220	672
CCT ATC AGA TCC AGT TAT TTC ACT TTT CAG CTT CAA AAT ATA GTT AAA Pro Ile Arg Ser Ser Tyr Phe Thr Phe Gln Leu Gln Asn Ile Val Lys 225 230 235	720
CCT TTG CCG CCA GTC TAT CTT ACT TTT ACT CGG GAG AGT TCA TGT GAA Pro Leu Pro Pro Val Tyr Leu Thr Phe Thr Arg Glu Ser Ser Cys Glu 240 245 250	768
ATT AAG CTG AAA TGG GGC ATA CCT TTG GGA CCT ATT CCA GCA AGG TGT Ile Lys Leu Lys Trp Gly Ile Pro Leu Gly Pro Ile Pro Ala Arg Cys 255 260 265	816
TTT GAT TAT GAA ATT GAG ATC AGA GAA GAT GAT ACT ACC TTG GTG ACT Phe Asp Tyr Glu Ile Glu Ile Arg Glu Asp Asp Thr Thr Leu Val Thr 270 275 280 285	864
GCT ACA GTT GAA AAT GAA ACA TAC ACC TTG AAA ACA ACA AAT GAA ACC Ala Thr Val Glu Asn Glu Thr Tyr Thr Leu Lys Thr Thr Asn Glu Thr 290 295 300	912
CGA CAA TTA TGC TTT GTA GTA AGA AGC AAA GTG AAT ATT TAT TGC TCA Arg Gln Leu Cys Phe Val Val Arg Ser Lys Val Asn Ile Tyr Cys Ser 305 310 315	960
GAT GAC GGA ATT TGG AGT GAG TGG AGT GAT AAA CAA TGC TGG GAA GGT Asp Asp Gly Ile Trp Ser Glu Trp Ser Asp Lys Gln Cys Trp Glu Gly 320 325 330	1008
GAA GAC CTA TCG AAG AAA ACT TTG CTA CGT TTC TGG CTA CCA TTT GGT Glu Asp Leu Ser Lys Lys Thr Leu Leu Arg Phe Trp Leu Pro Phe Gly 335 340 345	1056

TTC ATC TTA ATA TTA GTT ATA TTT GTA ACC GGT CTG CTT TTG CGT AAG 1104
 Phe Ile Leu Ile Leu Val Ile Phe Val Thr Gly Leu Leu Leu Arg Lys
 350 355 360 365

CCA AAC ACC TAC CCA AAA ATG ATT CCA GAA TTT TTC TGT GAT ACA TGAAGACTTT
 1159
 Pro Asn Thr Tyr Pro Lys Met Ile Pro Glu Phe Phe Cys Asp Thr
 370 375 380

CCTCTAGA 1167

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Ala Phe Val Cys Leu Ala Ile Gly Cys Leu Tyr Thr Phe Leu Ile
 1 5 10 15
 Ser Thr Thr Phe Gly Cys Thr Ser Ser Ser Asp Thr Glu Ile Lys Val
 20 25 30
 Asn Pro Pro Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr Leu Gly Tyr
 35 40 45
 Leu Tyr Leu Gln Trp Gln Pro Pro Leu Ser Leu Asp His Phe Lys Glu
 50 55 60
 Tyr Thr Val Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly Ser Glu Thr
 65 70 75 80
 Trp Lys Thr Ile Ile Thr Lys Asn Leu His Tyr Lys Asp Gly Phe Asp
 85 90 95
 Leu Asn Lys Gly Ile Glu Ala Lys Ile His Thr Leu Leu Pro Trp Gln
 100 105 110

Cys Thr Asn Gly Ser Glu Val Gln Ser Ser Trp Ala Glu Thr Thr Tyr
 115 120 125
 Trp Ile Ser Pro Gln Gly Ile Pro Glu Thr Lys Val Gln Asp Met Asp
 130 135 140
 Cys Val Tyr Tyr Asn Trp Gln Tyr Leu Leu Cys Ser Trp Lys Pro Gly
 145 150 155 160
 Ile Gly Val Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr Trp Tyr Glu
 165 170 175
 Gly Leu Asp Leu Ala Leu Gln Cys Val Asp Tyr Ile Lys Ala Asp Gly
 180 185 190
 Gln Asn Ile Gly Cys Arg Phe Pro Tyr Leu Glu Ala Ser Asp Tyr Lys
 195 200 205
 Asp Phe Tyr Ile Cys Val Asn Gly Ser Ser Glu Asn Lys Pro Ile Arg
 210 215 220
 Ser Ser Tyr Phe Thr Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro
 225 230 235 240
 Pro Val Tyr Leu Thr Phe Thr Arg Glu Ser Ser Cys Glu Ile Lys Leu
 245 250 255
 Lys Trp Gly Ile Pro Leu Gly Pro Ile Pro Ala Arg Cys Phe Asp Tyr
 260 265 270
 Glu Ile Glu Ile Arg Glu Asp Asp Thr Thr Leu Val Thr Ala Thr Val
 275 280 285
 Glu Asn Glu Thr Tyr Thr Leu Lys Thr Thr Asn Glu Thr Arg Gln Leu
 290 295 300
 Cys Phe Val Val Arg Ser Lys Val Asn Ile Tyr Cys Ser Asp Asp Gly
 305 310 315 320
 Ile Trp Ser Glu Trp Ser Asp Lys Gln Cys Trp Glu Gly Glu Asp Leu
 325 330 335
 Ser Lys Lys Thr Leu Leu Arg Phe Trp Leu Pro Phe Gly Phe Ile Leu
 340 345 350

Ile Leu Val Ile Phe Val Thr Gly Leu Leu Leu Arg Lys Pro Asn Thr
 355 360 365

Tyr Pro Lys Met Ile Pro Glu Phe Phe Cys Asp Thr
 370 375 380

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Trp Ser Xaa Trp Ser
 1 5

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1126 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 11..1126

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ACTTGGAGAA ATG GCT TTC GTC TAC TTG GCT ATC AGA TGC TTA TGT ACC
 Met Ala Phe Val Tyr Leu Ala Ile Arg Cys Leu Cys Thr
 1 5 10

TTT CTG ATA AGC ACA ACA TTC GGC TAT ACT TCA ACT TCA GAC ACC GAG	97
Phe Leu Ile Ser Thr Thr Phe Gly Tyr Thr Ser Thr Ser Asp Thr Glu	
15 20 25	
ATA AAA GTT AAC CCA CCT CAG GAT TTT GAG ATA GTG GAT CCC GGA TAT	145
Ile Lys Val Asn Pro Pro Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr	
30 35 40 45	
TTA GGT TAT CTC TAT TTG CAA TGG CAA CCC CCA CTG TCT CTG GAT AAT	193
Leu Gly Tyr Leu Tyr Leu Gln Trp Gln Pro Pro Leu Ser Leu Asp Asn	
50 55 60	
TTT AAG GAA TGC ACA GTG GAA TAT GAA CTA AAA TAC CGA AAC ATT GGT	241
Phe Lys Glu Cys Thr Val Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly	
65 70 75	
AGT GAA ACA TGG ACG ACC ATC ATT ACT AAG AAT CTA CAT TAC AAA GAT	289
Ser Glu Thr Trp Thr Thr Ile Ile Thr Lys Asn Leu His Tyr Lys Asp	
80 85 90	
GGG TTT GAT CTT AAC AAG GGC ATT GAA GCG AAG ATA CAC ACA CTT TTA	337
Gly Phe Asp Leu Asn Lys Gly Ile Glu Ala Lys Ile His Thr Leu Leu	
95 100 105	
CCA TGG CAA TGC ACA AAT GGA TCA GAA GTT CAA AGT TCC TGG GCA GAA	385
Pro Trp Gln Cys Thr Asn Gly Ser Glu Val Gln Ser Ser Trp Ala Glu	
110 115 120 125	
GCT ACT TAT TGG ATA TCG CCA CAA GGA ATT CCA GAA ACT AAA GTT CAG	433
Ala Thr Tyr Trp Ile Ser Pro Gln Gly Ile Pro Glu Thr Lys Val Gln	
130 135 140	
GAT ATG GAT TGT GTA TAT TAC AAT TGG CAA TAT TTA CTC TGT TCT TGG	481
Asp Met Asp Cys Val Tyr Tyr Asn Trp Gln Tyr Leu Leu Cys Ser Trp	
145 150 155	
AAA CCT GGC ATA GGT GTA CTT CTT GAT ACC AAT TAC AAC TTG TTT TAC	529
Lys Pro Gly Ile Gly Val Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr	
160 165 170	
TGG TAT GAG GGC TTG GAT CGT GCA TTA CAG TGT GTT GAT TAC ATC AAG	577
Trp Tyr Glu Gly Leu Asp Arg Ala Leu Gln Cys Val Asp Tyr Ile Lys	
175 180 185	

GTT GAT GGA CAA AAT ATT GGA TGC AGA TTT CCC TAT TTG GAG TCA TCA Val Asp Gly Gln Asn Ile Gly Cys Arg Phe Pro Tyr Leu Glu Ser Ser 190 195 200 205	625
GAC TAT AAA GAT TTC TAC ATT TGT GTT AAT GGA TCA TCA GAA ACC AAG Asp Tyr Lys Asp Phe Tyr Ile Cys Val Asn Gly Ser Ser Glu Thr Lys 210 215 220	673
CCT ATC AGA TCC AGT TAT TTC ACT TTT CAG CTT CAA AAT ATA GTT AAA Pro Ile Arg Ser Ser Tyr Phe Thr Phe Gln Leu Gln Asn Ile Val Lys 225 230 235	721
CCT TTG CCA CCA GTC TGT CTT ACT TGT ACT CAG GAG AGT TTA TAT GAA Pro Leu Pro Pro Val Cys Leu Thr Cys Thr Gln Glu Ser Leu Tyr Glu 240 245 250	769
ATT AAG CTG AAA TGG AGC ATA CCT TTG GGA CCT ATT CCA GCA AGG TGT Ile Lys Leu Lys Trp Ser Ile Pro Leu Gly Pro Ile Pro Ala Arg Cys 255 260 265	817
TTT GTT TAT GAA ATT GAG ATC AGA GAA GAT GAT ACT ACC TTG GTG ACT Phe Val Tyr Glu Ile Glu Ile Arg Glu Asp Asp Thr Thr Leu Val Thr 270 275 280 285	865
ACC ACA GTT GAA AAT GAA ACG TAC ACC TTG AAA ATA ACA AAT GAA ACC Thr Thr Val Glu Asn Glu Thr Tyr Thr Leu Lys Ile Thr Asn Glu Thr 290 295 300	913
CGA CAG TTA TGC TTT GTA GTA AGA AGC AAA GTG AAT ATT TAT TGC TCA Arg Gln Leu Cys Phe Val Val Arg Ser Lys Val Asn Ile Tyr Cys Ser 305 310 315	961
GAT GAT GGA ATT TGG AGT GAG TGG AGT GAT AAA CAA TGT TGG GAA GTT Asp Asp Gly Ile Trp Ser Glu Trp Ser Asp Lys Gln Cys Trp Glu Val 320 325 330	1009
GAA GAA CTA TTG AAG AAA ACT TTG CTA CTT TTC TTG TTA CCA TTT GGT Glu Glu Leu Leu Lys Lys Thr Leu Leu Leu Phe Leu Leu Pro Phe Gly 335 340 345	1057
TTC ATA TTA ATA TTA GTT ATA TTT GTA ACC GGT CTG CTT TTG TGT AAG Phe Ile Leu Ile Leu Val Ile Phe Val Thr Gly Leu Leu Leu Cys Lys 350 355 360 365	1105

AGA GAC AGC TAC CCG AAA ATG
 Arg Asp Ser Tyr Pro Lys Met
 370

1126

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 372 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Ala	Phe	Val	Tyr	Leu	Ala	Ile	Arg	Cys	Leu	Cys	Thr	Phe	Leu	Ile	1	5	10	15
Ser	Thr	Thr	Phe	Gly	Tyr	Thr	Ser	Thr	Ser	Asp	Thr	Glu	Ile	Lys	Val	20	25	30	
Asn	Pro	Pro	Gln	Asp	Phe	Glu	Ile	Val	Asp	Pro	Gly	Tyr	Leu	Gly	Tyr	35	40	45	
Leu	Tyr	Leu	Gln	Trp	Gln	Pro	Pro	Leu	Ser	Leu	Asp	Asn	Phe	Lys	Glu	50	55	60	
Cys	Thr	Val	Glu	Tyr	Glu	Leu	Lys	Tyr	Arg	Asn	Ile	Gly	Ser	Glu	Thr	65	70	75	80
Trp	Thr	Thr	Ile	Ile	Thr	Lys	Asn	Leu	His	Tyr	Lys	Asp	Gly	Phe	Asp	85	90	95	
Leu	Asn	Lys	Gly	Ile	Glu	Ala	Lys	Ile	His	Thr	Leu	Leu	Pro	Trp	Gln	100	105	110	
Cys	Thr	Asn	Gly	Ser	Glu	Val	Gln	Ser	Ser	Trp	Ala	Glu	Ala	Thr	Tyr	115	120	125	
Trp	Ile	Ser	Pro	Gln	Gly	Ile	Pro	Glu	Thr	Lys	Val	Gln	Asp	Met	Asp	130	135	140	
Cys	Val	Tyr	Tyr	Asn	Trp	Gln	Tyr	Leu	Leu	Cys	Ser	Trp	Lys	Pro	Gly	145	150	155	160

Ile Gly Val Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr Trp Tyr Glu
165 170 175

Gly Leu Asp Arg Ala Leu Gln Cys Val Asp Tyr Ile Lys Val Asp Gly
180 185 190

Gln Asn Ile Gly Cys Arg Phe Pro Tyr Leu Glu Ser Ser Asp Tyr Lys
195 200 205

Asp Phe Tyr Ile Cys Val Asn Gly Ser Ser Glu Thr Lys Pro Ile Arg
210 215 220

Ser Ser Tyr Phe Thr Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro
225 230 235 240

Pro Val Cys Leu Thr Cys Thr Gln Glu Ser Leu Tyr Glu Ile Lys Leu
245 250 255

Lys Trp Ser Ile Pro Leu Gly Pro Ile Pro Ala Arg Cys Phe Val Tyr
260 265 270

Glu Ile Glu Ile Arg Glu Asp Asp Thr Thr Leu Val Thr Thr Val
275 280 285

Glu Asn Glu Thr Tyr Thr Leu Lys Ile Thr Asn Glu Thr Arg Gln Leu
290 295 300

Cys Phe Val Val Arg Ser Lys Val Asn Ile Tyr Cys Ser Asp Asp Gly
305 310 315 320

Ile Trp Ser Glu Trp Ser Asp Lys Gln Cys Trp Glu Val Glu Glu Leu
325 330 335

Leu Lys Lys Thr Leu Leu Leu Phe Leu Leu Pro Phe Gly Phe Ile Leu
340 345 350

Ile Leu Val Ile Phe Val Thr Gly Leu Leu Leu Cys Lys Arg Asp Ser
355 360 365

Tyr Pro Lys Met
370

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9801

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TGGTCCTTCC CATGTTTCAC TACCA

25

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9941

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TTTCGGTATT TTAGTTCATA TTCCA

25

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9803

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGGAATTTGG AGTGAGTGGA GTGAT

25

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG9937

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TGAAGACCTA TCGAAGAAAA CTTTG

25

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG9800

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ATGGCTTTTCG TTTGCTTGGC TATCG

25

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZG9802

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CTCTTGATAT GGAAAGTCTT CATGTATC

28

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: AP1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CCATCCTAAT ACGACTCACT ATAGGGC

27

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: AP2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ACTCACTATA GGGCTCGAGC GGC

23

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9850

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TCTGATAGGC TTGTTCTCTG

20

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9851

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ATAGCCAAGC AAACGAAAGC

20

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9852

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ACCTGGCATA GGTGTACTTC

20

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG9919

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TTGCCGCCAG TCTATCTTAC

20

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG10317

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGGGGGTCTA GAGGAAAGTC TTCATGTATC ACAG

34

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 base pairs

(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG10319

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GGGGGGCTGG AGCTCGGAGA AATGGCTTTC GTT

33

- (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9820

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ACCCCCACTG TCTCTGGATC ATTTT

25

- (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG9806

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CACCTTCCA GCATTGTTTA TCACT

25

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG10320

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GGGGGGAGAT CTTAGACAC CGAGATAAAA GTT

33

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: ZG10318

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GGGGGGCTCG AGTTTCTTCG ATAGGTCTTC ACC

33

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:

(B) CLONE: ZG9882

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

TTACTCTGTT CTTGGAAACC TGG

23

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG10082

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ACTCTGTTCT TGGAAACCTG G

21

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG10083

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

AAATGAAACA TACACCTTGA AAAC

24

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZG10081

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GCATTGTTTA TCACTCCACT C

21

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZG9881

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TTCAC TTTGC TTCTTACTAC AAA

23

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZG10389

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GACTAGCAGA TCTGGGCTCT TTCTTCGATA GGTCTTCAC

39

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG10314

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TCGTGATTCT CTGGTCGGTG

20

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZG10315

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GTGATTGCTT TGGCGGTGAG

20

Claims

We claim:

1. An isolated polynucleotide encoding a ligand-binding receptor polypeptide, said polypeptide comprising a sequence of amino acids selected from the group consisting of:

- (a) residues 141 to 337 of SEQ ID NO:2;
- (b) allelic variants of (a); and
- (c) sequences that are at least 80% identical to (a) or (b).

2. An isolated polypeptide according to claim 1 comprising residues 141 to 337 of SEQ ID NO:2 or SEQ ID NO:4.

3. An isolated polynucleotide according to claim 1 wherein said polypeptide further comprises a transmembrane domain.

4. An isolated polynucleotide according to claim 3 wherein said transmembrane domain comprises residues 340 to 363 of SEQ ID NO:2, or an allelic variant thereof.

5. An isolated polynucleotide according to claim 3 wherein said polypeptide further comprises an intracellular domain.

6. An isolated polynucleotide according to claim 5 wherein said intracellular domain comprises residues 365 to 380 of SEQ ID NO:2, or an allelic variant thereof.

7. An isolated polynucleotide according to claim 1 wherein said polypeptide comprises residues 25 to 337 of SEQ ID NO:2 or SEQ ID NO:4.

8. An isolated polynucleotide according to claim 1 wherein said polypeptide comprises residues 1 to 380 of SEQ ID NO:2 or SEQ ID NO:4.

9. An isolated polynucleotide according to claim 1 which is a DNA as shown in SEQ ID NO:1 from nucleotide 49 to nucleotide 1188 or SEQ ID NO:3 from nucleotide 10 to nucleotide 1149.

10. An isolated polynucleotide according to claim 1 wherein said polypeptide further comprises an affinity tag.

11. An isolated polynucleotide according to claim 10 wherein said affinity tag is polyhistidine, protein A, glutathione S transferase, substance P, or an immunoglobulin heavy chain constant region.

12. An isolated polynucleotide according to claim 1 wherein said polynucleotide is DNA.

13. An expression vector comprising the following operably linked elements:

a transcription promoter;

a DNA segment encoding a secretory peptide and a ligand-binding receptor polypeptide, said polypeptide comprising a sequence of amino acids selected from the group consisting of:

(a) residues 141 to 337 of SEQ ID NO:2;

(b) allelic variants of (a); and

(c) sequences that are at least 80% identical to (a) or (b); and

a transcription terminator.

14. An expression vector according to claim 13 wherein said polypeptide comprises residues 141 to 337 of SEQ ID NO:2 or SEQ ID NO:4.

15. An expression vector according to claim 13 wherein said polypeptide further comprises a transmembrane domain.

16. An expression vector according to claim 15 wherein said transmembrane domain comprises residues 340 to 363 of SEQ ID NO:2, or an allelic variant thereof.

17. An expression vector according to claim 15 wherein said polypeptide further comprises an intracellular domain.

18. An expression vector according to claim 17 wherein said intracellular domain comprises residues 364 to 380 of SEQ ID NO:2, or an allelic variant thereof.

19. An expression vector according to claim 13 wherein said polypeptide comprises residues 25 to 337 of SEQ ID NO:2 or SEQ ID NO:4.

20. An expression vector according to claim 13 wherein said polypeptide comprises residues 1 to 380 of SEQ ID NO:2 or SEQ ID NO:4.

21. An expression vector comprising the following operably linked elements:

- (a) a transcription promoter;
- (b) a DNA segment encoding a secretory peptide and a chimeric polypeptide, wherein said chimeric polypeptide consists essentially of a first portion and a second portion joined by a peptide bond, said first portion consisting essentially of a ligand binding domain of a receptor polypeptide selected from the group consisting of:
 - (i) a receptor polypeptide as shown in SEQ ID NO:2;
 - (ii) allelic variants of SEQ ID NO:2; and

(iii) receptor polypeptides that are at least 80% identical to (i) or (ii),
and said second portion consisting essentially of an affinity tag; and

(c) a transcription terminator.

22. An expression vector according to claim 21 wherein said affinity tag is an immunoglobulin F_C polypeptide.

23. A cultured eukaryotic cell into which has been introduced an expression vector according to claim 13, wherein said cell expresses a receptor polypeptide encoded by the DNA segment.

24. A cell according to claim 23 wherein said cell further expresses a hematopoietic receptor β_C subunit.

25. A cell according to claim 23 wherein said cell is dependent upon an exogenously supplied hematopoietic growth factor for proliferation.

26. An isolated polypeptide comprising a segment selected from the group consisting of:

(a) residues 141 to 337 of SEQ ID NO:2;

(b) allelic variants of (a); and

(c) sequences that are at least 80% identical to (a) or (b),

wherein said polypeptide is substantially free of transmembrane and intracellular domains ordinarily associated with hematopoietic receptors.

27. A polypeptide according to claim 26 further comprising an immunoglobulin F_C polypeptide.

28. A polypeptide according to claim 26 further comprising an affinity tag.

29. A polypeptide according to claim 28 wherein said affinity tag is polyhistidine, protein A, glutathione S transferase, substance P, or an immunoglobulin heavy chain constant region.

30. A polypeptide according to claim 26 that is immobilized on a solid support.

31. A chimeric polypeptide consisting essentially of a first portion and a second portion joined by a peptide bond, said first portion consisting essentially of a ligand binding domain of a receptor polypeptide selected from the group consisting of:

- (a) a receptor polypeptide as shown in SEQ ID NO:2;
 - (b) allelic variants of SEQ ID NO:2; and
 - (c) receptor polypeptides that are at least 80% identical to (a) or (b),
- and said second portion consisting essentially of an affinity tag.

32. A polypeptide according to claim 31 wherein said affinity tag is an immunoglobulin F_C polypeptide.

33. A method for detecting a ligand within a test sample, comprising contacting a test sample with a polypeptide comprising a segment selected from the group consisting of:

- (a) residues 141 to 337 of SEQ ID NO:2;
 - (b) allelic variants of (a); and
 - (c) sequences that are at least 80% identical to (a) or (b),
- and detecting binding of said polypeptide to ligand in the sample.

34. A method according to claim 33 wherein said polypeptide comprises residues 25 to 337 of SEQ ID NO:2 or an allelic variant of SEQ ID NO:2.

35. A method according to claim 33 wherein said polypeptide further comprises transmembrane and intracellular domains.

36. A method according to claim 35 wherein said polypeptide is membrane bound within a cultured cell, and said detecting step comprises measuring a biological response in said cultured cell.



37. A method according to claim 36 wherein said biological response is cell proliferation or activation of transcription of a reporter gene.

38. A method according to claim 33 wherein said polypeptide is immobilized on a solid support.

39. An antibody that specifically binds to a polypeptide of claim 26.

1/1

Figure



INTERNATIONAL SEARCH REPORT

Internat. Application No.
PCT/US 97/04043

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K14/715 C07K19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94 29458 A (AMGEN INC) 22 December 1994 see the whole document	1-38
Y	WO 94 19463 A (AUCKLAND UNISERVICES LTD) 1 September 1994 see abstract see page 9, line 27-31 - page 10, line 1-21 see page 12, line 1-33	1-38
A	ENDOCRINOLOGY, vol. 127, no. 1, 1990, SPRINGFIELD, ILL, US, pages 251-258, XP002035642 T. TAKAO ET AL.: "Identification of interleukin-1 receptors in mouse testis" see the whole document	1-9, 13-20, 23,26, 33-37

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

22 July 1997

Date of mailing of the international search report

08.08.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/US 97/04043

Continuation DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 46, 1994, BETHESDA, MA,US, pages 29094-29101, XP002035641 J.W. BAUMGARTNER ET AL.,: "The role of the WSXWS equivalent motif in growth hormone receptor" see specially introduction and discussion see the whole document ---	1-9, 12-20, 26,33-37
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 87, 1990, WASHINGTON DC, US, pages 9655-9659, XP002035643 HAYASHIDA ET AL.,: "Molecular cloning of a second subunit of the receptor for human granulocyte-macrophage colony-stimulating factor (GM-CSF): reconstitution of a high affinity GM-CSF receptor" cited in the application see the whole document ---	1-9,26
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 93, 1 January 1996, WASHINGTON DC, US, pages 497-501, XP002010896 D.J. HILTON ET AL.: "Cloning and characterization of a binding subunit of the interleukin 13 receptor that is also a component of the interleukin 4 receptor" cited in the application see the whole document ---	1-9,26
A	WO 95 21930 A (ZYMOGENETICS INC) 17 August 1995 cited in the application see abstract ---	25
A	EP 0 488 170 A (BEHRINGWERKE AG) 3 June 1992 see abstract ---	26-30, 33,38
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 28, 1 July 1996, BETHESDA, MD,US, pages 16921-16926, XP002025819 D. CAPUT ET AL.,: "Cloning and characterization of a specific interleukin (11)-13 binding protein structurally related to the IL-5 receptor alpha chain" see specially figure 3 see the whole document ---	1-9,26

-/--

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/US 97/04043

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>THE JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, vol. 99, no. 1, 1 January 1997, ST. LOUIS, MO. US, page 287 XP000197588 M. COLLINS ET AL.,: "Comparison of two distinct ligand binding chains for the Il13 receptor in the mouse" see abstract</p> <p>-----</p>	1-9,26

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat. Application No

PCT/US 97/04043

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9429458 A	22-12-94	AU 7203694 A EP 0705341 A JP 8508889 T US 5521295 A	03-01-95 10-04-96 24-09-96 28-05-96
WO 9419463 A	01-09-94	AU 6117294 A	14-09-94
WO 9521930 A	17-08-95	US 5541085 A AU 1747195 A AU 1843595 A AU 7481094 A CA 2169173 A CA 2169190 A CA 2183266 A EP 0745125 A EP 0745132 A EP 0723456 A FI 960930 A FI 963185 A HU 75359 A NO 963371 A PL 315914 A WO 9521920 A WO 9521626 A	30-07-96 29-08-95 29-08-95 29-08-95 17-08-95 17-08-95 17-08-95 04-12-96 04-12-96 31-07-96 28-02-96 14-08-96 28-05-97 11-10-96 09-12-96 17-08-95 17-08-95
EP 0488170 A	03-06-92	DE 4037837 A AT 140084 T AU 660950 B AU 8814191 A CA 2056279 A DE 59107974 D ES 2090211 T JP 6160395 A US 5639597 A	04-06-92 15-07-96 13-07-95 04-06-92 29-05-92 08-08-96 16-10-96 07-06-94 17-06-97